

Seminar

Ramona Vogt

Lawrence Livermore Nat. Lab. and Univ. of California
at Davis

Heavy Flavor Azimuthal Correlations in Cold Nuclear Matter

It has been proposed that the azimuthal distributions of heavy flavor quark-antiquark pairs may be modified in the medium of a heavy-ion collision. This work tests this proposition through next-to-leading order (NLO) calculations of the azimuthal distribution, $d\sigma/d\phi$, including transverse momentum broadening, employing $\langle k_T^2 \rangle$ and fragmentation in exclusive $Q\bar{Q}$ pair production. While these studies were done for $p+p$, $p+\bar{p}$ and $p+\text{Pb}$ collisions, understanding azimuthal angle correlations between heavy quarks in these smaller, colder systems is important for their interpretation in heavy-ion collisions.

First, single inclusive p_T distributions calculated with the exclusive HVQMNR code are compared to those calculated in the fixed-order next-to-leading logarithm approach. Next the azimuthal distributions are calculated and sensitivities to $\langle k_T^2 \rangle$, p_T cut, and rapidity are studied at $\sqrt{s} = 7$ TeV. Finally, calculations are compared to $Q\bar{Q}$ data in elementary $p+p$ and $p+\bar{p}$ collisions at $\sqrt{s} = 7$ TeV and 1.96 TeV as well as to the nuclear modification factor $R_{p\text{Pb}}(p_T)$ in $p+\text{Pb}$ collisions at $\sqrt{s_{NN}} = 5.02$ TeV measured by ALICE.

The low p_T ($p_T < 10$ GeV) azimuthal distributions are very sensitive to the k_T broadening and rather insensitive to the fragmentation function. The NLO contributions can result in an enhancement at $\phi \sim 0$ absent any other effects. Agreement with the data was found to be good. The NLO calculations, assuming collinear factorization and introducing k_T broadening, result in significant modifications of the azimuthal distribution at low p_T which must be taken into account in calculations of these distributions in heavy-ion collisions.

Tuesday, 12.02.2019, 14:15 Uhr

Place: D6-135